

MoviPrep Split Dose Preparation for a Colonoscopy

COLONOSCOPY:

In order to achieve an accurate evaluation of your lower gastrointestinal (GI) tract, it is important that you properly prepare for the procedure. Doctors are able to obtain the clearest view of the bowel when it is flushed completely free of waste material. The thoroughness of this intestinal cleaning depends on you. Without your cooperation, the examination cannot accomplish its purpose and may have to be repeated.

Five (5) Days Before Procedure:

- DO NOT DRINK ALCOHOL FOR (5) DAYS PRIOR TO YOUR PROCEDURE.
- DO NOT take any aspirin or ibuprofen products. Tylenol (acetaminophen) is allowed.
- Stop taking anticoagulants or blood thinners such as: Coumadin, Naprosyn, Nuprin, Persantine, Plavix, Relafen, Rufen.

Day Before Procedure:

- You will need to follow a clear liquid/high sodium diet throughout the day. Items include: clear juices without pulp, clear soda, chicken or beef broth, popsicles, Jell-O, tea, coffee, etc. If you are diabetic, drink regular (non-diet) liquids in moderation throughout the day. Avoid dairy, juice products and dark colored liquids with red, purple, and blue dye.
- **STEP 1:** At ____ PM empty one Pouch A and one Pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. If preferred, you may mix the solution ahead of time and refrigerate prior to drinking. Use within 24 hours.
- **STEP 2:** The MoviPrep container is divided by 4 marks. Every 15 min, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed. Drink 16 oz of clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
- You may continue to have clear liquids until bedtime.
- DO NOT EAT OR DRINK ANYTHING **AFTER MIDNIGHT**.
- See attached sheet for further information.

Day of Procedure:

- Take any currently prescribed heart or blood pressure medications with a sip of water. If you are diabetic: To avoid dangerously low blood sugar, **Do Not** take your diabetic medication (pills or insulin) the morning of your procedure. You can resume your diabetes medications after the procedure once you have eaten.
- You may have clear liquids until 7:00 AM **ONLY IF YOUR PROCEDURE IS SCHEDULED AFTER 1:30 PM**.
- Plan on being at the facility for approximately 2-3 hours.
- A driver must accompany you to drive you home after your procedure. You will be sedated for your colonoscopy and will be drowsy for several hours after the procedure. You may not drive or work for 12 hours after the procedure.
- YOU WILL **NOT** BE ALLOWED TO DRIVE OR TAKE A CAB.
- **STEP 1:** At ____ AM empty one Pouch A and one Pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. If preferred, you may mix the solution ahead of time and refrigerate prior to drinking. Use within 24 hours.
- **STEP 2:** The MoviPrep container is divided by 4 marks. Every 15 min, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed. Drink 16 oz. of clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
- See attached sheet for further information.

Procedure Date: _____

Check-In Time: _____

Location: _____

MEDICAL CENTER OF PLANO


3901 W. 15th Street - Plano, TX 75075

(Enter thru ER entrance and go to the Out Patient Registration Desk)

PRESTON CROSSING ENDOSCOPY CENTER

3500 Preston Road, Suite 200 - Plano, TX 75093

1600 Coit Road, Suite 301 | Plano, Texas 75075 | 972.867.0019 | MatthewEidem.com



Preston Crossing Endoscopy Center
 3500 Preston Road
 Suite 200
 Plano, TX 75093
 Phone: 972-985-2300
 Fax: 972-985-2320